

# On the quasisymmetrical classification of infinitely renormalizable maps

## II. REMARKS ON MAPS WITH A BOUNDED TYPE TOPOLOGY

by Yunping Jiang

June, 1991

### §0 Introduction

This note is a remark to the paper [1]. The aim is to show that the techniques in [1] can also be used to understand the quasisymmetrical classification of infinitely renormalizable maps of bounded type. We will use the same terms and notations as those in [1] without further notices. The result we will prove is the following theorem.

**THEOREM 1.** *Suppose  $f$  and  $g$  in  $\mathcal{U}$  are two infinitely renormalizable maps of bounded type and topologically conjugate. Moreover, suppose  $H$  is the homeomorphism between  $f$  and  $g$ . Then  $H$  is quasisymmetric.*


Since the techniques as well as ideas of the proof are similar to those in [1], we outline the proof in the next section. The reader may refer to [1] and [2] for more details.

### §1 The outline of the proof of Theorem 1

We outline the proof of Theorem 1 by several lemmas.


Suppose  $f = h \circ Q_t$ , for some  $t > 1$ , in  $\mathcal{U}$  is an infinitely renormalizable map of bounded type. We note that  $Q_t(x) = -|x|^t$ . Let  $f_0 = f$ . And inductively, let  $f_k = \alpha_k \circ f_{k-1}^{n_k} \circ \alpha_k^{-1}$  be the renormalization  $\mathcal{R}(f_{k-1})$  of  $f_{k-1}$  where  $\alpha_k$  is the linear rescale from  $J_k$  to  $[-1, 1]$  and  $n_k$  is the return time for any integer  $k \geq 1$  (see [1]). We call  $J_k$  a restricted interval.

Let  $I_0$  be the interval  $[-1, 1]$  and  $I_k$  be the preimage of  $[-1, 1]$  under  $\alpha_1 \circ \cdots \circ \alpha_k$  for  $k \geq 1$ . We note that the set  $\{I_k\}_{k=0}^\infty$  forms a sequence of nested intervals. Moreover, one of the endpoints of  $I_k$ , say  $p_k$ , is a periodic point of period  $m_k = n_1 \cdots n_k$  of  $f$  and the orbit  $O(p_k)$  of  $p_k$  under  $f$  stays outside of the interior of  $I_k$  (see Figure 1).



**Figure 1**

Suppose  $L_k$  is the image of  $I_k$  under  $f^{\circ m_k}$  and  $T_k$  is the interval bounded by the points  $p_k$  and  $p_{k+1}$ . Let  $M_k$  be the complement of  $T_k$  in  $L_k$ . Then  $M_k$  is the interval bounded by  $p_{k+1}$  and  $c_{m_k}$ , where  $c_{m_k} = f^{\circ m_k}(0)$  (See Figure 2).



**Figure 2**

LEMMA 1. *There is a constant  $C_1 = C_1(f) > 0$  such that*

$$C_1^{-1} \leq |M_k|/|I_k| \leq C_1.$$

*for all the integers  $k \geq 0$ .*

*Proof.* This lemma is actually proved in [3] by using the techniques such as the smallest interval and shuffle permutation on the intervals.

LEMMA 2. *There is a constant  $C_2 = C_2(f) > 0$  such that*

$$C_2^{-1} \leq |I_k|/|I_{k-1}| \leq C_2$$

*for all the integers  $k \geq 0$ .*

We first prove a more general result, as that in [1], as follows. Let  $\mathcal{K} = \mathcal{K}(t, N, K)$ , for fixed numbers  $t > 1$ ,  $N \geq 2$  and  $K > 0$ , be the subspace of renormalizable maps  $f = h \circ Q_t$  in  $\mathcal{U}$  such that  $|(N(h))(x)| \leq K$  for all  $x$  in  $[-1, 0]$  and all the return times  $n_k$  of  $\mathcal{R}^{\circ k}(f)$  are less than or equal to  $N$ .

LEMMA 3. *There is a constant  $C_3 = C_3(t, N, K) > 0$  such that*

$$C_3^{-1} \leq f(0) = c_1(f) \leq C_3$$

*for all  $f$  in  $\mathcal{K}$ .*

*Proof.* The proof of this lemma is similar to the proof of Lemma 3 in [1] but needs little more work to solve a little more complicated equation.

Remember that  $f_k$  is the  $k^{th}$ -renormalization of  $f = f_0$ . Let  $f_k = h_k \circ Q_t$ . We note that the graph of  $f_k$  is the rescale of the graph of the restriction of  $f^{o m_k}$  to  $I_k$ .

LEMMA 4. *There is a constant  $C_4 = C_4(f) > 0$  such that*


$$|(N(h_k))(x)| \leq C_4$$

for all  $x$  in  $[-1, 0]$  and all the integer  $k \geq 0$ .

*Proof.* It is the a prior bound proved in [3].

*Proof of Lemma 2.* It is now a direct corollary of Lemma 1, Lemma 3 and Lemma 4 for  $K = C_4$  and  $N = \max_{0 \leq k < \infty} \{n_k\}$ .

The set of the nested intervals  $\{I_0, I_1, \dots, I_k, \dots\}$  gives a partition of  $[-1, 1]$  as follows. Let  $p_k$  be one of the endpoints of  $I_k$  and  $O_{k,f}(p_k)$  be the intersection of  $I_{k-1}$  and the orbit  $O_{k,f}(p_k)$  of  $p_k$  under  $f$  for  $k \geq 1$ . Suppose  $M_{k-1,1}, \dots, M_{k-1,n_k+1}$  are the connected components of  $I_{k-1} \setminus (O_{k,f}(p_k) \cup I_k)$  for  $k \geq 1$ . Then the set  $\eta_0 = \{M_{k,i}\}$  for  $i = 1, \dots, n_k + 1$  and  $k = 1, 2, \dots$  forms a partition of  $[-1, 1]$  (see Figure 3).



**Figure 3**


Now we are going to define a Markov map  $F$  induced by  $f$ . Let  $F$  be a function of  $[-1, 1]$  defined by

$$F(x) = \begin{cases} f(x), & x \in M_{1,1} \cup M_{1,2} \cup \dots \cup M_{1,n_1+1}, \\ f^{o n_1}(x), & x \in M_{2,1} \cup \dots \cup M_{2,n_2+1}, \\ \vdots \\ f^{o n_1 n_2 \dots n_k}(x), & x \in M_{k,1} \cup \dots \cup M_{k,n_k+1}, \\ \vdots \end{cases}$$

It is clearly that  $F$  is a Markov map in the sense that the image of every  $M_{k,i}$  under  $F$  is the union of some intervals in  $\eta_0$  (Figure 4).

Let  $g_{k,i} = (F|_{M_{k,i}})^{-1}$  for  $k = 1, \dots$ , and  $i = 1, \dots, n_k + 1$  be the inverse branches of  $F$  with respect to the Markov partition  $\eta_0$ . Suppose  $w = i_0 i_1 \dots i_{l-1}$  is a finite sequence of the set  $\mathcal{I} = \{(k, i), k = 1, \dots \text{ and } i = 1, \dots, n_k + 1\}$ . We say it is admissible if the range  $M_{i_s}$  of  $g_{i_s}$  is contained in the domain  $F_{i_{s-1}}(J_{i_{s-1}})$  of  $g_{i_{s-1}}$  for  $s = 1, \dots, l - 1$ . For an admissible

sequence  $w = i_0 i_1 \cdots i_{l-1}$ , we can define  $g_w = g_{i_0} \circ g_{i_1} \circ \cdots \circ g_{i_{l-1}}$ . We use  $D(g_w)$  to denote the domain of  $g_w$  and use  $|D(g_w)|$  to denote the length of the interval  $D(g_w)$ .



**Figure 4**

**DEFINITION 1.** We say the induced Markov map  $F$  has bounded distortion property if there is a constant  $C_5 = C_5(f) > 0$  such that

- (a)  $C_5^{-1} \leq |M_{k,i}|/|M_{k,i+1}| \leq C_5$  for  $k = 1, 2, \dots$  and  $i = 1, \dots, n_k$ ,
- (b)  $C_5^{-1} \leq |M_{k,i}|/|I_k| \leq C_5$  for  $k = 1, 2, \dots$  and  $i = 1, \dots, n_k + 1$ , and
- (b)  $|(N(g_w))(x)| \leq C_5/|D(g_w)|$  for all  $x$  in  $D(g_w)$  and all admissible  $w$ .

The reason we give this definition is the following lemma as that in [1].

**LEMMA 5.** Suppose  $f$  and  $g$  in  $\mathcal{U}$  are two infinitely renormalizable maps of bounded type and  $H$  is the conjugacy between  $f$  and  $g$ . If both of the induced Markov maps  $F$  and  $G$  have the bounded distortion property, then  $H$  is quasimetric.

*Proof.* It can be proved by almost the same arguments as that we used in the paper [2]. For more details of the proof, the reader may refer to [2].


Now the proof of Theorem 1 concentrates on the next lemma.

**LEMMA 6.** Suppose  $f = h \circ Q_t$ , for some  $t > 1$ , in  $\mathcal{U}$  is an infinitely renormalizable map of bounded type and  $F$  is the Markov map induced by  $f$ . Then  $F$  has the bounded distortion

property.

*Proof.* Let  $I_{k,j} = \left(f^{\circ m_{k-1}}|I_{k-1}\right)^{\circ j}(I_k)$  for  $j = 0, 1, \dots, n_k$  and  $\{G_{k,i}\}$  are all the connected components of  $I_k \setminus \cup_{j=0}^{n_k} I_{k,j}$  (Figure 5).

Each  $M_{k,j}$  is either a single  $G_{k,i}$  or  $I_{k,j} \cup G_{k,i}$  for some  $j$  and some  $i$ . By the bounded geometry [3] of  $\{I_{k,j}\}$  and  $\{G_{k,i}\}$ , there is a constant  $C_6 > 1$  such that all the ratios  $|I_{k,j}|/|I_{k,j'}|$ ,  $|G_{k,i}|/|G_{k,i'}|$  and  $|G_{k,i}|/|I_{k,j}|$  are in the interval  $[C_6^{-1}, C_6]$ . We note that  $C_6$  does not depend on  $k$  as well as  $i, i', j$  and  $j'$ . This fact and Lemma 3 imply the condition (a) in Definition 1.



**Figure 5**

The condition (b) in Definition 1 is assured by Lemma 2 and the condition (a). The proof of the condition (c) in Definition 1 is similar to that in [1].

The arguments in Lemma 1 to Lemma 6 give the proof of Theorem 1.

## References

- [1] Y. Jiang, On quasimetric classification of infinitely renormalizable maps – Maps with Feigenbaum's topology, preprint in this issue, IMS at SUNY at Stony Brook.
- [2] Y. Jiang, Dynamics of certain smooth one-dimensional mappings – II. Geometrically finite one-dimensional mappings, preprint 1991/1, IMS, SUNY at Stony Brook.
- [3] D. Sullivan, Bounds, quadratic differentials, and renormalization conjectures, preprint, 1991 and American Mathematical Society Centennial Publications, Volume 2: Mathematics into the Twenty-first Century, to appear.

Yunping Jiang  
Institute for Mathematical Sciences  
SUNY at Stony Brook  
Stony Brook, NY 11794  
e-mail: jiang@math.sunysb.edu